For Research Use Only

CoraLite® Plus 555 Anti-Human CD25/IL-2RA (BC96)

Catalog Number: CL555-65096



Purification Method:

Affinity purification

Recommended Dilutions:

FC: 5 ul per 10^6 cells in 100 μ l

Excitation/Emission maxima

CloneNo.:

suspension

wavelengths:

554 nm / 570 nm

BC96

Basic Information

Catalog Number: CL555-65096 Concentration: 100tests, 5 μl/test

Source: Mouse Isotype:

IgG1, kappa

GenBank Accession Number: BC156627 GeneID (NCBI):

UNIPROT ID: P01589 Full Name: interleukin 2 receptor, alpha

Calculated MW: 272 aa, 31 kDa

Positive Controls:

FC: PHA treated human PBMCs,

Applications

Tested Applications: FC

Species Specificity:

human

Background Information


 $Proliferation of \ T\ lymphocytes\ is\ triggered\ by\ the\ interaction\ of\ IL-2\ with\ its\ specific\ receptor\ following\ T$ lymphocyte activation. The receptor for IL-2 has three forms, generated by different combinations of three different proteins, the alpha chain (IL-2R alpha), the beta chain (IL-2R beta), and the gamma chain (IL-2R gamma) (PMID: 8476561). IL-2R alpha (also known as CD25) is a type I transmembrane protein present on activated T cells, activated B cells, some thymocytes, myeloid precursors, and oligodendrocytes. CD25 is also found on natural CD4+Foxp3+ Treg cells and a subset of human CD4+ memory T cells (PMID: 22585674).

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data



1x10^6 PHA treated (3 days) human PBMCs were surface stained with 5 ul CoraLite® Plus 555 Anti-Human CD25 (CL555-65096, Clone:BC96) (red) or CoraLite® Plus 555 Mouse IgG1 Isotype Control (blue). 1x10^6 untreated human PBMCs were surface stained with 5 ul CoraLite® Plus 555 Anti-Human CD25 (CL555-65096, Clone:BC96) (black dashed). Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Lymphocytes were gated.



1x10^6 PHA treated (3 days) human PBMCs were surface stained with CL750 Anti-Human CD3 and 5 ul Coralite® Plus 555 Anti-Human CD25 (CL555-65096, Clone:BC96) (right) or Coralite® Plus 555 Mouse IgG1 Isotype Control (left). 1x10^6 untreated human PBMCs were surface stained with 5 ul Coralite® Plus 555 Anti-Human CD25 (CL555-65096, Clone:BC96) (center). Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Lymphocytes were gated.