For Research Use Only

CoraLite®555-conjugated IFITM2/3 Monoclonal antibody



Catalog Number: CL555-66081

Featured Product

Basic Information

Catalog Number: CL555-66081 Concentration:

1000 μg/ml Source: Mouse

lgG1 Immunogen Catalog Number:

AG17863

Isotype:

GenBank Accession Number:

BC070243 GeneID (NCBI): 10410

interferon induced transmembrane

protein 3 (1-8U) Observed MW:

15-20 kDa

Full Name:

Purification Method:

Protein G purification

CloneNo.: 2E8D12

Recommended Dilutions: IF/ICC: 1:50-1:500

FC (Intra): 0.40 ug per 10^6 cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 557 nm / 570 nm

Applications

Tested Applications: IF/ICC, FC (Intra) Species Specificity:

human

Positive Controls:

IF/ICC: U2OS cells, FC (Intra): HeLa cells,

Background Information

IFITM3, also named as interferon-inducible protein 1-8U, belongs to the CD225 family. It is IFN-induced antiviral protein that mediates cellular innate immunity to at least three major human pathogens, namely influenza A H1N1 virus, West Nile virus (WNV), and dengue virus, by inhibiting the early steps of replication. IFITM3 is identified as $interferon-induced\ cellular\ proteins\ that\ restrict\ infections\ by\ retroviruses\ and\ filoviruses\ and\ of\ influenza\ virus\ of\ influenza\ of\ i$ flaviviruses, respectively. IFITM3, the most potent antiviral IFITM, was found to inhibit an uncharacterized early infectious event after VSV endocytosis, but before primary transcription of its viral genome. IFITM proteins are viral restriction factors that can inhibit infection mediated by the influenza A virus (IAV) hemagglutinin (HA) protein. They differentially restrict the entry of a broad range of enveloped viruses, and modulate cellular tropism independently of viral receptor expression. This antibody recognizes both IFITM2 and IFITM3.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed U2OS cells using Coralite®555 IFITM2/3 antibody (CL555-66081, Clone: 2E8D12) at dilution of 1:200.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite®555 Anti-Human IFITM2/3 (CL555-66081, Clone:2E8D12) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).