For Research Use Only

## CoraLite®594-conjugated Beta Tubulin Polyclonal antibody

www.ptglab.com

Catalog Number: CL594-10068

**Basic Information** 

Catalog Number: CL594-10068 Concentration:

1000 ug/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG0117

GenBank Accession Number:

BC000748 GeneID (NCBI): 10381 **UNIPROT ID:** Q13509 Full Name:

tubulin, beta 3 Calculated MW: 450 aa, 50 kDa

Observed MW: 50-55 kDa

**Purification Method:** 

Antigen affinity purification Recommended Dilutions:

IF/ICC: 1:200-1:500

FC (Intra): 0.80 ug per 10^6 cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

**Applications** 

**Tested Applications:** IF/ICC, FC (Intra) Species Specificity: human

Positive Controls: IF/ICC: HeLa cells, FC (Intra): HeLa cells,

**Background Information** 

There are five tubulins in human cells: alpha, beta, gamma, delta, and epsilon. Tubulins are conserved across species. They form heterodimers, which multimerize to form a microtubule filament. An alpha and beta tubulin heterodimer is the basic structural unit of microtubules. The heterodimer does not come apart, once formed. The alpha and beta tubulins, which are each about 55 kDa MW, are homologous but not identical. Alpha, beta, and gamma tubulins have all been used as loading controls. Tubulin expression may vary according to resistance to antimicrobial and antimitotic drugs.


Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data







Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CoraLite®594 Beta Tubulin antibody (CL594-10068) at dilution of 1:200.