### For Research Use Only

# CoraLite®594-conjugated GOLGA2/GM130 Polyclonal antibody



Catalog Number: CL594-11308

Featured Product

1 Publications

**Basic Information** 

Catalog Number: CL594-11308 Concentration: 1000 µ g/ml Source:

Rabbit Isotype:

Immunogen Catalog Number:

AG1848

AG1040

GenBank Accession Number:

BC014188
GeneID (NCBI):
2801
UNIPROT ID:
Q08379
Full Name:

golgi autoantigen, golgin subfamily

a, 2

Calculated MW: 111 kDa Purification Method:

Antigen affinity purification Recommended Dilutions:

IF/ICC: 1:50-1:500 FC (Intra): 0.20 ug per 10^6 cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

**Applications** 

Tested Applications: IF/ICC, FC (Intra)
Cited Applications:

IF

Species Specificity:

human Cited Species: human Positive Controls:

IF/ICC : HepG2 cells,
FC (Intra) : HepG2 cells,

## **Background Information**

GOLGA2, also known as GM130, is a 130 kDa cis-Golgi matrix protein, one component of the detergent and salt-resistant Golgi matrix. It is a peripheral membrane protein highly bound to the Golgi membrane and localized mainly at the cytoplasmic face of the cis-Golgi membrane. Together with its interacting partner proteins, including p115, giantin, GRASP65, and Rab GTPase, GOLGA2/GM130 is involved in regulating ER-to-Golgi transport and maintaining the Golgi structure. Emerging evidence suggests that the GOLGA2/GM130 has potential roles in the control of glycosylation, cell cycle progression, and higher-order cell functions such as cell polarization and directed cell migration. (PMID: 20197635)

#### **Notable Publications**

Qiu Pan 38253527 Nat Commun IF	Author	Pubmed ID	Journal	Application
	Qiu Pan	38253527	Nat Commun	IF

## Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

# Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite®594 GOLGA2/GM130 antibody (CL594-11308) at dilution of 1:200.



1X10^6 HepG2 cells were intracellularly stained with 0.2 ug CoraLite® 594 Anti-Human GOLGA2/GM130 (CL594-11308) (red), or 0.2 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).