For Research Use Only

CoraLite®594-conjugated iNOS Polyclonal antibody

Catalog Number: CL594-18985

Featured Product

4 Publications



Basic Information

Catalog Number: CL594-18985 Concentration: 1000 μg/ml Source:

Rabbit Isotype:

Observed MW: 110-130 kDa, 65-70 kDa

131 kDa

GenBank Accession Number:

NM_000625 GeneID (NCBI): 4843 **UNIPROT ID:** P35228 Full Name:

nitric oxide synthase 2, inducible Calculated MW:

Purification Method:

Antigen affinity purification Recommended Dilutions:

IF/ICC: 1:50-1:500

FC (Intra): 0.80 ug per 10⁶ cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

Applications

Tested Applications: IF/ICC, FC (Intra) Cited Applications: IF

Species Specificity: human, mouse, rat Cited Species: mouse

Positive Controls:

IF/ICC: HeLa cells, FC (Intra): HepG2 cells,

Background Information

NOS2, also named as iNOS and NOS2A, produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body. NO is a reactive free radical which acts as a biologic mediator in several processes, including neurotransmission, antimicrobial and antitumoral activities. NOS2 is a nitric oxide synthase which is expressed in liver and is inducible by a combination of lipopolysaccharide and certain cytokines. iNOS has a very short half-life due to rapid degradation by calpain. iNOS monomer is a direct substrate of calpain I and can be cleaved by calpain I at the canonical CaM-binding site(503-532aa) of iNOS, and then a ~70kD band can be detected by western (PMID:11786228). This antibody is specific to NOS2.

Notable Publications

Author	Pubmed ID	Journal	Application
Rongxin Wan	39894624	Biomater Adv	IF
Banxin Luo	39724782	Mol Immunol	IF
Ganghua Yang	37827053	Biomaterials	IF


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1X10^6 HepG2 cells were intracellularly stained with 0.8 ug Coralite®594 Anti-Human iNOS (CL594-18985) (red), or 0.8 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using CoraLite®594 iNOS antibody (CL594-18985) at dilution of 1:200.