For Research Use Only

# CoraLite®594-conjugated Caspase 3/P17/P19 Polyclonal antibody



**Purification Method:** 

IF/ICC: 1:50-1:500

100 µl suspension

Antigen affinity purification

Excitation/Emission maxima

FC (Intra): 0.40 ug per 10<sup>6</sup> cells in a

Recommended Dilutions:

Catalog Number: CL594-19677

Featured Product

**Basic Information** 

Catalog Number: CL594-19677 Concentration: 1000 ug/ml

Source: Rabbit Isotype:

Calculated MW: 32 kDa Observed MW:

32-35 kDa, 17 kDa, 19 kDa

GenBank Accession Number:

NM 004346 GeneID (NCBI):

P42574 Full Name:

**UNIPROT ID:** 

wavelengths: caspase 3, apoptosis-related cysteine 588 nm / 604 nm peptidase

**Applications** 

**Tested Applications:** IF/ICC, FC (Intra) Species Specificity: human, mouse, rat

Positive Controls:

IF/ICC: NIH/3T3 cells, FC (Intra): HepG2 cells,

#### **Background Information**

Caspases, a family of endoproteases, are critical players in cell regulatory networks controlling inflammation and cell death. Initiator caspases (caspase-2, -8, -9, -10, -11, and -12) cleave and activate downstream effector caspases (caspase-3, -6, and -7), which in turn execute apoptosis by cleaving targeted cellular proteins. Caspase 3 (also named CPP32, SCA-1, and Apopain) proteolytically cleaves poly(ADP-ribose) polymerase (PARP) at the beginning of apoptosis. Caspase 3 plays a key role in the activation of sterol regulatory element binding proteins (SREBPs) between the basic helix-loop-helix leucine zipper domain and the membrane attachment domain. Caspase 3 can also form a heterocomplex with other proteins and performs the molecular mass of 50-70 kDa. This antibody can recognize p17, p19, and p32 of Caspase 3.

### Storage


#### Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage


## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed NIH/3T3 cells using CoraLite®594 Caspase 3/p17/p19 antibody (CL594-19677) at dilution of 1:100.



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug CoraLite®594 Anti-Human Caspase 3/p17/p19 (CL594-19677) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using CoraLite® 594 Caspase 3/p17/p19 antibody (CL594-19677) at dilution of 1:200.