For Research Use Only

CoraLite®594-conjugated NeuN Polyclonal antibody



Catalog Number: CL594-26975 1 Publications

Basic Information

Catalog Number: GenBank Accession Number:
CL594-26975 NM_001082575
Concentration: GeneID (NCBI):
1000 ug/ml 146713
Source: UNIPROT ID:

Rabbit A6NFN3
Isotype: Full Name:

Immunogen Catalog Number: Observed MW:

AG25689 46-52 kDa

Purification Method:
Antigen affinity purification
Recommended Dilutions:

IF-P: 1:50-1:500

FC (Intra): 0.40 ug per 10^6 cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

Applications

Tested Applications: IF-P, FC (Intra) Cited Applications: IF

Species Specificity: human, mouse, rat, pig Cited Species:

rat

Positive Controls:

IF-P: mouse brain tissue, rat brain tissue

FC (Intra): U-87 MG cells,

Background Information

NeuN, encoded by FOX3, is a neuron-specific nuclear protein. Anti-NeuN stains exclusively neuronal cells in the central and peripheral nervous systems, especially postmitotic and differentiating neurons, as well as terminally differentiated neurons. Anti-NeuN has been used widely as a reliable tool to detect most postmitotic neuronal cell types. The immunohistochemical staining is primarily localized in the nucleus of the neurons with lighter staining in the cytoplasm. Several isoforms of NeuN exist due to the alternative splicing. Although the predicted MW of NeuN are 34/35 kDa, it was detected as doublet around 46-52 kDa. (PMID: 21747913)

hexaribonucleotide binding protein 3

Notable Publications

Author	Pubmed ID	Journal	Application
Beibei Liu	39415414	Mol Pain	IF

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using CoraLite®594 NeuN antibody (CL594-26975) at dilution of 1:200.



1X10^6 U-87 MG cells were intracellularly stained with 0.4 ug Coralite®594 Anti-Human NeuN (CL594-26975) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded rat brain tissue using CoraLite®594 NeuN antibody (CL594-26975) at dilution of 1:200, CoraLite® Plus 488 GFAP antibody (CL488-60190, Clone: 4B2E10, green). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded rat brain tissue using CoraLite®594 NeuN antibody (CL594-26975) at dilution of 1:200, CoraLite® Plus 488 GFAP antibody (CL488-60190, Clone: 4B2E10, green). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).