For Research Use Only

CoraLite® Plus 594 Anti-Human CD163 (GHI/61)

Antibodies | ELISA kits | Proteins www.ptglab.com

Catalog Number: CL594-65169

Basic Information

Catalog Number: CL594-65169

Concentration: 100tests, 5 ul/test

Source: Mouse Isotype:

IgG1, kappa

1156 a

Applications Tested Applications:

Species Specificity:

human

GenBank Accession Number:

BC051281 GeneID (NCBI):

9332 ENSEMBL Gene ID: ENSG00000177575

UNIPROT ID: Q86VB7 Full Name:

CD163 molecule
Calculated MW:
1156 aa, 125 kDa

Purification Method:
Protein G purification

Protein G purification

CloneNo.: GHI/61

Recommended Dilutions: FC: 5 ul per 10^6 cells in 100 $\,\mu$ l

suspension

Positive Controls:

FC: human PBMCs,

Excitation/Emission maxima

wavelengths: 594 nm / 615 nm

Background Information

CD163, also known as M130, is a membrane glycoprotein which belongs to the scavenger receptor superfamily (PMID: 8370408). It is an acute phase-regulated and signal-inducing macrophage protein expressed exclusively in monocytes and tissue macrophages (PMID: 11196644). CD163 mediates endocytosis of haptoglobin-haemoglobin complexes (PMID: 11196644). The uptake of haptoglobin by macrophages contributes to the recycling of iron and also to the inflammatory response (PMID: 22900885). Soluble CD163 (sCD163), as a result of ectodomain shedding during inflammatory activation of macrophages, circulates in blood and has been suggested as a plasma/serum marker for macrophage activity (PMID: 12570164).


Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:


PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data





1x10^6 human PBMCs were surface stained with CL488 Anti-Human CD14 (CL488-65246) and 5 ul CoraLite®594 Anti-Human CD163 (CL594-65169, Clone:GHI/61) or CoraLite®594 Mouse IgG1 Isotype Control (CL594-65124, Clone: MOPC-21). Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Monocytes were gated.



1x10^6 human PBMCs were surface stained with 5 ul CoraLite®594 Anti-Human CD163 (CL594-65169, Clone:GHI/61) or CoraLite®594 Mouse IgG1 Isotype Control (CL594-65124, Clone: MOPC-21). Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Monocytes were gated.