For Research Use Only

# CoraLite®594-conjugated HDAC1 Monoclonal antibody

Catalog Number: CL594-66085



**Purification Method:** 

CloneNo.:

4C4G7

Protein G purification

Recommended Dilutions:

Excitation/Emission maxima

FC (Intra): 0.20 ug per 10^6 cells in a

IF/ICC: 1:4000-1:16000

100 µl suspension

wavelengths:

588 nm / 604 nm

**Basic Information** 

Catalog Number: GenBank Accession Number: CL594-66085 BC000301

Concentration: GeneID (NCBI): 3065

Source: UNIPROT ID:
Mouse Q13547
Isotype: Full Name:

lgG1 histone deacetylase 1

Immunogen Catalog Number:Calculated MW:AG1912855 kDa

Observed MW: 60-65 kDa

**Applications** 

Tested Applications: IF/ICC, FC (Intra) Species Specificity: human, mouse, rat Positive Controls:

IF/ICC : A431 cells, HeLa cells
FC (Intra) : HeLa cells,

## **Background Information**

Histone deacetylases(HDAC) are a class of enzymes that remove the acetyl groups from the lysine residues leading to the formation of a condensed and transcriptionally silenced chromatin. The protein encoded by this gene belongs to the histone deacetylase/acuc/apha family and is a component of the histone deacetylase complex, which is responsible for gene expression silencing. It also plays an important role in the control of cell proliferation and differentiation by interacting with RB, p53 and other transcription factors. At least 4 classes of HDAC were identified. As a class I HDAC, HDAC 1 was primarily found in the nucleus. This antibody is raised against residues near the C terminus of human HDAC1. The calculated molecular weight of HDAC1 is 55 kDa, but modified HDAC1 is about 60-65 kDa.

### Storage


#### Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage


#### **Selected Validation Data**




Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CL594-66085 (HDAC1 antibody) at dilution of 1:100.



1X10^6 HeLa cells were intracellularly stained with 0.2 ug CoraLite®594 Anti-Human HDAC1 (CL594-66085, Clone:4C4G7) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (4% PFA) fixed A431 cells using Coralite®594 HDAC1 antibody (CL594-66085, Clone: 4C4G7) at dilution of 1:8000.



Immunofluorescent analysis of (4% PFA) fixed A431 cells using CoraLite®594 HDAC1 antibody (CL594-66085, Clone: 4C4G7) at dilution of 1:8000.