#### For Research Use Only

# CoraLite®594-conjugated TUBB3specific/TUJ1 Monoclonal antibody



Catalog Number: CL594-66375 2 Publications

**Basic Information** 

Catalog Number: CL594-66375

Concentration: 1000 ug/ml Source: Mouse

Isotype: IgG1 GenBank Accession Number:

NM\_001197181 GeneID (NCBI): 10381 Full Name:

tubulin, beta 3

Calculated MW:
55 kDa

Observed MW:

50-55 kDa

Purification Method:

Protein G purification

CloneNo.: 1F8G10

Recommended Dilutions:

WB: 1:1000-1:4000 IF-P: 1:50-1:500

FC (Intra): 0.40 ug per 10^6 cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

**Applications** 

Tested Applications: WB, IF-P, FC (Intra)

Cited Applications:

IHC, IF

Species Specificity: human, mouse, rat Cited Species: mouse Positive Controls:

WB: SH-SY5Y cells, PC-12 cells, Neuro-2a cells

IF-P: mouse brain tissue, FC (Intra): SH-SY5Y cells,

## **Background Information**

TUBB3, the class III  $\,eta\,$  tubulin or Tuj1, is selectively expressed in testis and neurons of the central and peripheral nervous system. It has been widely used as a marker for neurons. Aberrant expression of TUBB3 has also been found in various tumors of non-neural origin and can be used as a biomarker for cancer aggressiveness and a marker for the tendency to respond poorly to chemotherapy. This antibody is specific to TUBB3 but not cross-react with other tubulin isoforms. And the antibody is conjugated with CL594, Ex/Em 593 nm/614 nm.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Nao Iguchi	35332157	Sci Rep	IF
Yuki Shindo	35427599	Exp Cell Res	IHC

### Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3  $\,$ 

Aliquoting is unnecessary for -20°C storage


#### Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with CL594-66375 (TUBB3-specific/TUJ1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immun of luorescent analysis of (4% PFA) fixed mouse brain tissue using CL594-66375 (TUBB3-specific antibody) at dilution of 1:50.



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using CL594-66375 (TUBB3-specific antibody) at dilution of 1:50.



1X10^6 SH-SY5Y cells were intracellularly stained with 0.4 ug Coralite®594 Anti-Human TUBB3-specific (CL594-66375, Clone:1F8G10) (red), or 0.4 ug Mouse IgG1 Isotype Control (CL594-6636, Clone: TIF8D3F10) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using CoraLite®594 TUBB3-specific antibody (CL594-66375, Clone: 1F8G10 ) at dilution of 1:200.