For Research Use Only

## CoraLite® Plus 647-conjugated STAT3 Polyclonal antibody



Catalog Number: CL647-10253

Featured Product

**Basic Information** 

Catalog Number: CL647-10253 Concentration: 1000 µg/ml

Source: Rabbit Isotype:

Immunogen Catalog Number:

AG0360

GenBank Accession Number:

BC000627 GeneID (NCBI): 6774 UNIPROT ID: P40763 Full Name:

signal transducer and activator of transcription 3 (acute-phase response

factor)

Calculated MW: 770 aa, 88 kDa Observed MW: 88 kDa Purification Method:

Antigen affinity purification
Recommended Dilutions:

FC (Intra): 0.20 ug per 10^6 cells in a 100 µl suspension

Excitation/Emission maxima

wavelengths: 654 nm / 674 nm

**Applications** 

Tested Applications: FC (Intra)

Species Specificity: human, mouse, rat

Positive Controls:

FC (Intra): HeLa cells,

## **Background Information**

Signal transducer and activator of transcription 3 (acute-phase response factor) (STAT3, synonyms: APRF, FLJ20882, MGC16063) is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. STAT3 is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. STAT3 mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of STAT3. This antibody is a rabbit polyclonal antibody raised against residues near the N terminus of human STAT3. STAT3 exists three isoforms and the molecular weight of each isoform respectively is 83 kDa, 87 kDa and 88 kDa.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer

PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



1X10^6 HeLa cells were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human STAT3 (CL647-10253) (red), or 0.2 ug isotype control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).