For Research Use Only

CoraLite® Plus 647-conjugated c-MYC Polyclonal antibody



Catalog Number: CL647-10828

Featured Product

Basic Information

Catalog Number: CL647-10828 Concentration: 1000 μg/ml

Source: Rabbit Isotype:

Immunogen Catalog Number:

AG1263

oncogene homolog (avian) Calculated MW:

> 49 kDa Observed MW: 62-65 kDa, 50 kDa

GenBank Accession Number:

v-myc myelocytomatosis viral

BC000141

4609

P01106

GeneID (NCBI):

UNIPROT ID:

Full Name:

Purification Method:

Antigen affinity purification

Recommended Dilutions:

FC (Intra): 0.20 ug per 10^6 cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 654 nm / 674 nm

Applications

Tested Applications: FC (Intra)

Species Specificity:

human

Positive Controls:

FC (Intra): HeLa cells,

Background Information

MYC contains one basic helix-loop-helix (bHLH) domain. This protein is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. It seems to activate the transcription of growth-related genes. It binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. This antibody is a rabbit polyclonal antibody raised against recombinant protein of human MYC.The 50kDa band recognized by antibody is the native form of MYC, while the other bands, between 60-70kDa, are the phosphorylated form of MYC [PMID: 12189186]

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1X10^6 HeLa cells were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human c-MYC (CL647-10828) (red), or 0.2 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).