For Research Use Only

CoraLite® Plus 647-conjugated CDKN2A/P16-INK4A Polyclonal antibody



Purification Method:

100 µl suspension

wavelengths:

Antigen affinity purification

Excitation/Emission maxima

FC (Intra): 0.80 ug per 10^6 cells in a

Recommended Dilutions:

Catalog Number: CL647-10883

Featured Product

Basic Information

Catalog Number:
CL647-10883

Concentration:
1000 ug/ml
1029

Source:
Rabbit
P42771

Isotype:
GenBank Accession Number:
GeneID (NCBI):
1029
UNIPROT ID:
P42771
Full Name:

gG cyclin-dependent kinase inhibitor 2A 654 nm / 674 nm

Immunogen Catalog Number: Calculated MW: 16 kDa

Observed MW:

16-18 kDa

Applications

Tested Applications: FC (Intra)
Species Specificity:

human

Positive Controls: FC (Intra): HeLa cells,

Background Information

CDKN2A generates several transcript variants which differ in their first exons. At least three alternatively-spliced variants encoding distinct proteins proteins were reported. Two of them named p16-INK4 and p14 are sharing 50% identity. The third one -p14(ARF) is entirely unrelated. 10883-1-AP reacts with p16 specifically. P16 plays an essential role in regulating the cell cycle, and mutations in p16 increase the risk of developing various cancers, including melanoma.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1x10^6 HeLa cells were intracellularly stained with 0.8 ug CoraLite® Plus 647 Anti-Human P16-INK4A (CL647-10883)(red), or 0.8 ug CoraLite® Plus 647-conjugated Rabbit IgG control Rabbit PolyAb (CL647-30000) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).