For Research Use Only

CoraLite® Plus 647-conjugated HSP90 Monoclonal antibody

www.ptglab.com

Purification Method:

Protein A purification

Recommended Dilutions:

Excitation/Emission maxima

100 µl suspension

wavelengths:

654 nm / 674 nm

FC (Intra): 0.20 ug per 10^6 cells in a

CloneNo.:

3F11C1

Catalog Number: CL647-60318

Basic Information

Catalog Number: CL647-60318

Concentration: 1000 ug/ml Source: Mouse

IgG2a Immunogen Catalog Number:

AG3826

Isotype:

Tested Applications: FC (Intra)

Species Specificity: human, mouse, rat, pig GenBank Accession Number:

BC023006 GeneID (NCBI): 3320 **UNIPROT ID:** P07900

Full Name: heat shock protein 90kDa alpha (cytosolic), class A member 1

Calculated MW: 853 aa, 90 kDa Observed MW:

85-90 kDa

Positive Controls:

FC (Intra): K-562 cells,

Background Information

HSP90, encoded by HSP90AA1, is a constitutively and ubiquitously expressed molecular chaperone that is crucial for $the \ stability \ and \ function \ of \ many \ proteins. \ HSP90 \ provides \ chaperoning \ activity \ for \ client \ proteins; \ many \ of \ them$ are members of oncogenic pathways, indicating its implication in tumor malignancy. HSP90 mainly resides in the cytosol, while it can also be released to the extracellular space. Secreted Hsp90 is a C-terminal truncated form. It has been reported that the level of plasma Hsp90 is positively correlated with tumor malignancy in clinical cancer patients, and can be a promising diagnostic marker for tumor malignancy in clinical application. This antibody may cross-react with HSP90AB1 due to the high homology between them.

Storage


Applications

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1X10^6 K-562 cells were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human HSP90 (CL647-60318, Clone:3F11C1) (red), or 0.2 ug isotype control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).