

CoraLite® Plus 750-conjugated GAPDH Monoclonal antibody

Catalog Number: **CL750-60004**

Basic Information

Catalog Number:

CL750-60004

Size:

1000 µg/ml

Source:

Mouse

Isotype:

IgG2b

Immunogen Catalog Number:

AG0766

GenBank Accession Number:

BC004109

GeneID (NCBI):

2597

UNIPROT ID:

P04406

Full Name:glyceraldehyde-3-phosphate
dehydrogenase**Calculated MW:**

36 kDa

Observed MW:

36 kDa

Purification Method:

Protein A purification

CloneNo.:

1E6D9

Recommended Dilutions:

WB 1:2000-1:10000

**Excitation/Emission maxima
wavelengths:**

755 nm / 780 nm

Applications

Tested Applications:

WB

Species Specificity:

human, mouse, rat, yeast, plant, zebrafish

Positive Controls:

WB : Jurkat cells, HeLa cells

Background Information

Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) catalyzes the phosphorylation of glyceraldehyde-3-phosphate during glycolysis. GAPDH participates in nuclear events including transcription, binding RNA, RNA transportation, DNA replication, DNA repair and apoptosis. Being stably and constitutively expressed at high levels in most tissues and cells, GAPDH is considered a housekeeping protein. It is widely used as a control for RT-PCR and also loading control in electrophoresis and Western blotting. GAPDH is normally expressed in cellular cytoplasm or membrane, but can occasionally translocate to the nucleus after the addition of post-translational modifications such as S-nitrosylation. This antibody is raised against full length GAPDH of human origin. It can recognize the 36 kDa GAPDH protein in most cells/tissues. In addition, a band below 36 kDa can always be detected as the isoform or spliced product of GAPDH (PMID: 23885286, 23877755, 19368702). Please note that some physiological factors, such as hypoxia and diabetes, increase GAPDH expression in certain cell types. For murine tissue samples, conjugated mouse antibody HRP-60004 and rabbit antibody 10494-1-AP are preferable.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with CL750-60004 (GAPDH antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.