For Research Use Only

Cardinal Red™-conjugated P62/SQSTM1 Polyclonal antibody



Catalog Number: CR-18420

Featured Product

Basic Information

Catalog Number: CR-18420

Concentration: 1000 µg/ml

Source: Rabbit

Isotype:

Immunogen Catalog Number:

AG13131

48 kDa Observed MW:

62 kDa

BC017222

8878

Q13501

GeneID (NCBI):

UNIPROT ID:

Full Name:

sequestosome 1 Calculated MW:

GenBank Accession Number:

Purification Method:

Antigen affinity purification

Recommended Dilutions:

FC (Intra): 0.20 ug per 10^6 cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 592 nm / 611 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity:

human

Positive Controls:

FC (Intra): HeLa cells,

Background Information

Sequestosome 1 (SQSTM1/p62) is a multifunctional adaptor protein implicated in selective autophagy, cell signaling pathways, and tumorigenesis. p62 has been implicated in shuttling ubiquitinated and aggregated proteins for autophagic degradation. p62 is degraded during the autophagic process, which makes its intracellular level a $marker for autophagy\ progression.\ p62\ is\ at\ the\ cross-roads\ of\ several\ signaling\ pathways\ including\ Ras/\ Raf/\ MAPK$ and NF x B and plays important role in cancer. p62 is a component of inclusion bodies/ protein aggregates found in human diseases, including Huntington's disease, Alzheimer's disease, Parkinson's disease, and nephropathic cystinosis (PMID: 22074114, 22860231, 22714671). The molecular weight of p62 is predicted to be 48/38 kDa (depending on the isoform), while western blot analyses using this antibody detects the bands around 45-48 kDa and 60-62 kDa, respectively.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1X10^6 HeLa cells were intracellularly stained with 0.2 ug Cardinal Red™ Anti-Human P62,5Q5TM1 (CR-18420) (red), or 0.2 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).