For Research Use Only

Cardinal Red™ Anti-Human CD3 (UCHT1)

Catalog Number: CR-65151



Purification Method:

Affinity purification

Recommended Dilutions:

FC: 5 ul per 10^6 cells in 100 μ l

Excitation/Emission maxima

CloneNo.:

suspension

wavelengths:

592 nm / 611 nm

UCHT1

Basic Information

Catalog Number: CR-65151 Concentration:

100tests, 5 µ l/test Source: Mouse Isotype:

Isotype: IgG1, kappa

Tested Applications:

Species Specificity: Human GenBank Accession Number:

BC049847 GeneID (NCBI): 916

Calculated MW: 207 aa, 23 kDa

ENSEMBL Gene ID: ENSG00000198851 UNIPROT ID: P07766

Full Name: CD3e molecule, epsilon (CD3-TCR complex)

Positive Controls:

FC : human PBMCs,

Background Information

CD3 is a multimeric protein associated with the T-cell receptor (TCR) to form a complex involved in antigen recognition and signal transduction (PMID: 15885124). CD3 is composed of CD3 γ , δ , ϵ , and ζ chains (PMID: 1826255). It is expressed by thymocytes in a developmentally regulated manner, T cells, and some NK cells (PMID: 3289580). The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction (PMID: 11985657).

Storage


Applications

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer.

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data



1X10^6 human PBMCs were surface co-stained with Atlantic Blue Anti-Human CD19 and 5 ul Cardinal Red™ Anti-Human CD3 (CR-65151, Clone:UCHT1) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul Cardinal Red™ Anti-Human CD3 (CR-65151, Clone:UCHT1) (red) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.