## For Research Use Only

## Cardinal Red™ Anti-Human CD163 (GHI/61)

Catalog Number: CR-65169



**Basic Information** 

Catalog Number: GenBank Accession Number: CR-65169 BC051281

GeneID (NCBI): Concentration: 9332

100tests, 5 ul/test

ENSEMBL Gene ID: Source: Mouse ENSG00000177575 Isotype: UNIPROT ID: IgG1, kappa Q86VB7 Full Name:

> CD163 molecule Calculated MW:

1156 aa, 125 kDa

**Purification Method:** 

Purified by protein-A affinity

chromatography

CloneNo.: GHI/61

Recommended Dilutions:

FC: 5 ul per 10^6 cells in 100  $\mu$  l

suspension

Excitation/Emission maxima

wavelengths: 592 nm / 611 nm

**Applications** 

**Tested Applications:** 

Species Specificity:

Human

Positive Controls:

FC: human PBMCs,

## **Background Information**


CD163, also known as M130, is a membrane glycoprotein which belongs to the scavenger receptor superfamily (PMID: 8370408). It is an acute phase-regulated and signal-inducing macrophage protein expressed exclusively in monocytes and tissue macrophages (PMID: 11196644). CD163 mediates endocytosis of haptoglobin-haemoglobin  $complexes (PMID: 11196644). The \ uptake \ of \ haptoglobin \ by \ macrophages \ contributes \ to \ the \ recycling \ of \ iron \ and$ also to the inflammatory response (PMID: 22900885). Soluble CD163 (sCD163), as a result of ectodomain shedding during inflammatory activation of macrophages, circulates in blood and has been suggested as a plasma/serum marker for macrophage activity (PMID: 12570164).

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

## Selected Validation Data



1X10^6 human PBMCs were surface stained with 5 ul Cardinal Red™ Anti-Human CD163 (CR-65169, Clone:GHI/61) (red) or unstained. Cells were not fixed. Monocytes and granulocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul Cardinal Red™ Anti-Human CD163 (CR-65169, Clone:GHI/61) or unstained. Cells were not fixed. Monocytes and granulocytes were gated.