For Research Use Only

FITC Plus Anti-Human CD3 (UCHT1)

Catalog Number: FITC-65151



Basic Information

Catalog Number: FITC-65151 Concentration:

100tests, 5 $\,\mu$ l/test

Source: Mouse Isotype: IgG1, kappa

Isotype: IgG1, kappa

complex)

Calculated MW:
207 aa, 23 kDa

BC049847

916

GeneID (NCBI):

ENSEMBL Gene ID:

ENSG00000198851

UNIPROT ID:

Full Name:

P07766

GenBank Accession Number:

CD3e molecule, epsilon (CD3-TCR

Purification Method: Affinity purification

CloneNo.: UCHT1

Recommended Dilutions: FC: 5 ul per 10^6 cells in 100 $\,\mu$ l

suspension
Excitation/Emission maxima

wavelengths: 495 nm / 524 nm

Applications

Tested Applications:

Species Specificity:

Human

Positive Controls:

FC: human PBMCs, human peripheral blood

lymphocytes

Background Information

CD3 is a multimeric protein associated with the T-cell receptor (TCR) to form a complex involved in antigen recognition and signal transduction (PMID: 15885124). CD3 is composed of CD3 γ , δ , ϵ , and ζ chains (PMID: 1826255). It is expressed by thymocytes in a developmentally regulated manner, T cells, and some NK cells (PMID: 3289580). The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction (PMID: 11985657).


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer.

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data



1X10^6 human PBMCs were surface co-stained with CL647 Plus Anti-Human CD19 and 5 ul FITC Plus Anti-Human CD3 (FITC-65151, Clone:UCHT1) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated



1X10^6 human PBMCs were surface stained with 5 ul FITC Plus Anti-Human CD3 (FITC-65151, Clone:UCHT1) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated