

Catalog Number: HZ-1138

Data Sheet





Animal Component-Free


Human cell expressed

Tag-Free


Endotoxin Free

Product Description				
Animal-free Recombinant Human Activin A is a member of the TGF beta superfamily, expressed from human 293 cells as disulfide linked homodimer with an apparent molecular mass of 25 kDa. This cytokine is produced in a serum-free, chemically defined media.				
Alternative Names	Alternative Names Activin A, Activin beta A chain, EDF, FRP, INHBA, Inhibin beta A chain, inhibin, beta A			
Source Human Embryonic Kidney cells (HEK293). HEK293-derived Activin A protein				

Specifications					
Test	Method	Specification			
Activity	Dose-dependent inhibition of the proliferation of the MPC-11 cell line (mouse plasmocytoma cell line)	0.5-3.5 ng/mL			
Molecular Mass	SDS-PAGE	13kDa reduced, 22 kDa non-reduced, homodimer, non- glycosylated			
Purity	SDS-PAGE	>95%			
Endotoxin	LAL	<1 EU/ µ g			



Recombinant human Activin A (HZ-1138) causes dosedependent inhibition of proliferation in the Mouse plasmacytoma (MPC-11) cell line. Proliferation of the MPC-11 cell line was assessed using Promega CellTiter 96®. MPC-11 cells were treated with increasing concentrations of recombinant human Activin A for 72 hrs hours before addition of CellTiter96® reagent. The



www.ptglab.com

Document #: FR-QA118-101 Rev 0 Data Sheet Version #: 1 Proteintech Group, Inc.

5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com

Preparation				
Shipping Temperature	amnient temperature			
Formulation	1x PBS, See Certificate of Analysis for details			
Reconstitution Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein to 0.2 mg/mL in sterile 1x PBS p containing 0.1% endotoxin-free recombinant human serum albumin (HSA). Gently Swirl or tap vial to mix				

	Product Form	Temperature Conditions	Storage Time (From Date of Receipt)
	Lyophilized	-20°C to -80°C	Until Expiry Date
Stability and Storage	Lyophilized	Room Temperature	2 weeks
, -	Reconstituted as per CofA	-20°C to -80°C	6 months
	Reconstituted as per CofA	4°C	1 week
	Avoid repeated freeze-thaw cycles.		

www.ptglab.com

Document #: FR-QA118-101 Rev 0 Data Sheet Version #: 1 Proteintech Group, Inc.

5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com