# For Research Use Only

# PE Anti-Human CD163 (GHI/61)

Catalog Number: PE-65169 6 Publications





#### **Basic Information**

Catalog Number:
PE-65169
Concentration:
100tests, 5 ul/test

Source: Mouse Isotype: IgG1, kappa GenBank Accession Number:

BC051281 GeneID (NCBI): 9332 ENSEMBL Gene ID: ENSG00000177575

UNIPROT ID: Q86VB7 Full Name: CD163 molecule Calculated MW: 1156 aa, 125 kDa **Purification Method:** 

The purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography.

CloneNo.: GHI/61

Recommended Dilutions: FC: 5 ul per 10^6 cells in 100  $\,\mu$  l suspension

Excitation/Emission maxima wavelengths:

496 nm, 565 nm / 578 nm

## **Applications**

**Tested Applications:** 

FC

Cited Applications:

FC

Species Specificity:

Human
Cited Species:
human

#### Positive Controls:

FC : human PBMCs.

# **Background Information**

CD163, also known as M130, is a membrane glycoprotein which belongs to the scavenger receptor superfamily (PMID: 8370408). It is an acute phase-regulated and signal-inducing macrophage protein expressed exclusively in monocytes and tissue macrophages (PMID: 11196644). CD163 mediates endocytosis of haptoglobin-haemoglobin complexes (PMID: 11196644). The uptake of haptoglobin by macrophages contributes to the recycling of iron and also to the inflammatory response (PMID: 22900885). Soluble CD163 (sCD163), as a result of ectodomain shedding during inflammatory activation of macrophages, circulates in blood and has been suggested as a plasma/serum marker for macrophage activity (PMID: 12570164).

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Tao Yan	35410993	Cell Death Discov	FC
Yijun Lu	40184814	Int Immunopharmacol	FC
Tianci Han	39891717	Cancer Immunol Immunother	FC


### Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

# Selected Validation Data



1X10^6 human PBMCs were treated with FC receptor block then surface stained with CL405 Anti-Human CD14 and 5 ul PE Anti-Human CD163 (PE-65169, Clone:GHI/61) or PE Mouse IgG1 Isotype Control. Cells were not fixed. Monocytes were gated.



1X10^6 human PBMCs were treated with FC receptor block then surface stained with 5 ul PE Anti-Human CD163 (PE-65169, Clone:GHI/61) or PE Mouse IgG1 Isotype Control. Cells were not fixed. Monocytes were gated.