



产品介绍	该别藻蓝蛋白-链霉亲和素偶联物（APC Streptavidin）是荧光标记物（别藻蓝蛋白，APC）和链霉亲和素通过共价键结合的偶联产物，为溶液状态。建议每次测试使用量为 0.0125 ug/test ，务必每次实验都进行浓度优化以确保得到最佳染色，适用于生物素标记的抗体、核酸、蛋白或者其他生物素标记分子的检测。										
产品成分	<table><tr><th>名称</th><th>别藻蓝蛋白-链霉亲和素偶联物</th></tr><tr><td>浓度</td><td>0.2 mg/ml</td></tr><tr><td>存储溶液</td><td>PBS with 0.1% sodium azide and 0.5% BSA，pH 7.4.</td></tr><tr><td>激发波长</td><td>650 nm</td></tr><tr><td>发射波长</td><td>660 nm</td></tr></table>	名称	别藻蓝蛋白-链霉亲和素偶联物	浓度	0.2 mg/ml	存储溶液	PBS with 0.1% sodium azide and 0.5% BSA，pH 7.4.	激发波长	650 nm	发射波长	660 nm
名称	别藻蓝蛋白-链霉亲和素偶联物										
浓度	0.2 mg/ml										
存储溶液	PBS with 0.1% sodium azide and 0.5% BSA，pH 7.4.										
激发波长	650 nm										
发射波长	660 nm										
包装规格	100 ug/500 ug/1 mg										
保存条件	2°C-8°C避光保存，一年有效。										
注意	该别藻蓝蛋白-链霉亲和素偶联物无需稀释保存，2°C-8°C避光储存，不要冷冻。										

Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Biotin PEX14 antibody (Biotin-10594) at dilution of 1:200 and APC Streptavidin (Cat.No. PF00022), CL488-phalloidin (green).



Flow cytometric analysis of human PBMCs using Biotin anti-human CD3 and APC Streptavidin: human PBMCs were stained with Biotin CD3 Recombinant Antibody (Cat.No. Biotin-65569, Clone: OKT3), washed with PBS, and stained with APC Streptavidin (Cat.No. PF00022) and analyzed by flow cytometry.